

Probleme pentru pregătirea concursurilor

A. Nivel gimnazial

G186. Fie m, n, x, y numere naturale nenule astfel încât $m < n$, $\frac{m+n}{(m,n)}$ este număr par, iar $x^m = y^n$. Demonstrați că $x - y$ nu este pătrat perfect.

Geanina Hăvârneanu, Iași

G187. Considerăm numerele naturale a, b, c, d și x , astfel încât ac, bd și $ad + bc$ sunt divizibile cu x . Arătați că bc se divide cu x .

Ciprian Baghiu, Iași

G188. Determinați tripletele $(p, q, r) \in \mathbb{Z}^3$ pentru care

$$[a^{pq} + b^{pq} + (a^p + b^p)^q]^r = r[a^{pqr} + b^{pqr} + (a^p + b^p)^{qr}], \quad \forall a, b \in \mathbb{R}.$$

Temistocle Bîrsan, Iași

G189. Se consideră trei sfere având volumele v_1, v_2, v_3 , ariile s_1, s_2, s_3 și lungimile cercurilor ecuatoriale l_1, l_2, l_3 . Demonstrați că

$$\frac{v_1}{l_1 + l_2} + \frac{v_2}{l_2 + l_3} + \frac{v_3}{l_3 + l_1} \geq \frac{1}{12\pi}(s_1 + s_2 + s_3).$$

D.M. Bătinețu-Giurgiu, București

G190. Dacă x și y sunt numere iraționale astfel încât $x^2 + y, y^2 + x$ și $x + y$ sunt raționale, demonstrați că $xy < \frac{1}{4}$.

Neculai Moraru, Suceava și Silviu Boga, Iași

G191. Fie $A = \{1, 2, \dots, n\}$, unde $n \in \mathbb{N}, n \geq 5$. Demonstrați că A poate fi scrisă ca reuniunea a două mulțimi disjuncte B și C , astfel încât produsul elementelor lui B să fie egal cu suma elementelor lui C .

Gheorghe Iurea, Iași

G192. Demonstrați că în orice triunghi neisoscel ABC , există $M, N \in [BC]$, $P, Q \in [AB]$ și $R, S \in [AC]$ astfel încât segmentele $[AU], [BT]$ și $[CV]$ sunt laturi ale unui triunghi ascuțitunghic, oricare ar fi punctele $U \in [MN], V \in [PQ]$ și $T \in [RS]$.

Marius Drăgan, București

G193. Folosind efectiv 24 bețe de chibrit, construiți un poligon având aria de 7 unități, unde unitatea de arie este pătratul cu latura de un băț de chibrit. (*În legătură cu o problemă din "Amuzamente matematice" de Martin Gardner*)

Dumitru Mihalache, Bârlad

G194. Fie $ABCD$ un patrulater convex, iar semidreptele $[CK]$ și $[CL]$ sunt bisectoarele unghiurilor \widehat{ACD} , respectiv \widehat{ACB} ($K \in AD, L \in AB$). Demonstrați că dreapta LK trece prin centrul cercului înscris în triunghiul ABD dacă și numai dacă patrulaterul $ABCD$ este inscriptibil.

Neculai Roman, Mircești (Iași)

G195. Punctele M și P aparțin laturii neoparalele (AD) a trapezului $ABCD$, astfel încât $M \in [AP]$. Paralelele prin M și P la baze intersectează BC în N , respectiv Q . Demonstrați că trapezele $ABNM$ și $PQCD$ au diagonalele respectiv paralele dacă și numai dacă CD și PQ sunt direct proporționale cu MN și AB (*În legătură cu G105 din RecMat 1/2006.*)

Claudiu Ștefan Popa, Iași

B. Nivel liceal

L186. Determinați măsurile unghiurilor triunghiului ABC , știind că

$$\operatorname{tg} A + 2\sqrt{\operatorname{tg} B} + 3\sqrt[3]{\operatorname{tg} C} = 6\sqrt[6]{\operatorname{tg} A + \operatorname{tg} B + \operatorname{tg} C}.$$

Cătălin Calistru, Iași

L187. Se consideră triunghiul ABC și punctele $D, L \in (BC)$, $E, M \in (CA)$ și $F, N \in (AB)$. Notăm $\{P\} = FE \cap AL$, $\{Q\} = DF \cap BM$, $\{S\} = ED \cap CN$. Demonstrați că dacă două dintre următoarele afirmații sunt adevărate, atunci este adevărată și a treia:

- i) dreptele AD, BE și CF sunt concurente;
- ii) dreptele AL, BM și CN sunt concurente;
- iii) dreptele DP, EQ și FS sunt concurente.

Titu Zvonaru, Comănești

L188. Fie Oxy un reper cartezian, dreapta $d : x = 1$ și un punct $M \in d$. Punctul P se obține din M printr-o rotație de centru O și unghi 2α , unde α este unghiul orientat \widehat{yOM} . Scrieți ecuația $y = f(x)$ a curbei descrise de P atunci când M parcurge dreapta d și desenați graficul acestei curbe.

Temistocle Bîrsan, Iași

L189. Un sistem de $n \geq 3$ puncte coplanare A_1, A_2, \dots, A_n , oricare trei necoliniare, se numește *bun* atunci când oricare ar fi $1 \leq i < j < k \leq n$ și oricare ar fi punctele $M \in [A_i A_j]$, $N \in [A_j A_k]$ și $P \in [A_k A_i]$, are loc inegalitatea $MN^2 + NP^2 + MP^2 \leq A_i A_j^2 + A_j A_k^2 + A_k A_i^2$. Determinați sistemele bune cu număr maxim de puncte.

Vlad Emanuel, student, București

L190. Demonstrați că

$$8\sqrt{15} \leq (\sqrt{x} + \sqrt{x+48})(\sqrt{16-x} + \sqrt{1-x}) \leq 36, \quad \forall x \in [0, 1].$$

Gheorghe Iurea, Iași

L191. Arătați că, pentru orice numere reale pozitive x, y, z , are loc inegalitatea

$$\begin{aligned} (x^2 + y^2 + z^2) \left(\frac{1}{(x+y)(x+z)} + \frac{1}{(x+y)(y+z)} + \frac{1}{(x+z)(y+z)} \right) &\geq \frac{9}{4} \geq \\ &\geq (xy + xz + yz) \left(\frac{1}{(x+y)(x+z)} + \frac{1}{(x+y)(y+z)} + \frac{1}{(x+z)(y+z)} \right). \end{aligned}$$

Marian Tetiva, Bârlad

L192. Pe \mathcal{P} mulțimea polinoamelor unitare având coeficienții în intervalul $[0, 9]$. Determinați supremumul modulelor rădăcinilor complexe ale polinoamelor din \mathcal{P} .

Adrian Reisner, Paris

L193. Fie $A \in M_3(\mathbb{R})$ o matrice simetrică, având urma egală cu determinantul. Demonstrați că $2 \cdot \text{Tr}(A^{n+2}) \geq \text{Tr} A \cdot (\text{Tr}(A^{n+1}) - \text{Tr}(A^{n-1}))$, $\forall n \in \mathbb{N}^*$.

Paul Georgescu și Gabriel Popa, Iași

L194. Fie $p \geq 2$ un număr natural și ecuația $\frac{1}{\sqrt[p]{1+x}} + \frac{1}{\sqrt[p]{2+x}} + \dots + \frac{1}{\sqrt[p]{n+x}} = \sqrt[p]{x^{p-1}}$. Demonstrați că, pentru fiecare $n \in \mathbb{N}^*$, ecuația are o unică soluție pozitivă x_n , arătați că șirul $(x_n)_{n \geq 1}$ este monoton și calculați $\lim_{n \rightarrow \infty} \frac{n}{x_n}$.

Gabriel Dospinescu, student, Paris

L195. Fie $(a_n)_{n \geq 1}$ un șir de numere reale mai mari ca 1, astfel încât $a_{n+1} \geq 2a_n - 1$, $\forall n \in \mathbb{N}^*$. Definim $x_n = \frac{1}{a_1} + \frac{1}{a_1 a_2} + \dots + \frac{1}{a_1 a_2 \dots a_n}$, $\forall n \geq 1$.

a) Demonstrați că șirul $(x_n)_{n \geq 1}$ este convergent (notăm cu x limita sa).

b) Arătați că șirul $(\alpha_n)_{n \geq 1}$, unde $\alpha_n = x_n + \frac{\alpha_n}{a_1 a_2 \dots a_n (a_{n+1} - 1)}$, $\forall n \geq 1$, este monoton, convergent și determinați limita sa.

Dumitru Mihalache și Marian Tetiva, Bârlad

Training problems for mathematical contests

A. Junior highschool level

G186. Let m, n, x, y be four nonzero natural numbers such that $m < n$, $\frac{m+n}{(m,n)}$ is an even number, and $x^m = y^n$. Prove that $x - y$ is not a perfect square.

Geanina Hăvârneanu, Iași

G187. Let us consider the natural numbers a, b, c, d and x such that ac, bd and $ad + bc$ are divisible by x . Show that bc is divisible by x .

Ciprian Baghiu, Iași

G188. Determine the triples $(p, q, r) \in \mathbb{Z}^3$ such that

$$[a^{pq} + b^{pq} + (a^p + b^p)^q]^r = r[a^{pqr} + b^{pqr} + (a^p + b^p)^{qr}], \forall a, b \in \mathbb{R}.$$

Temistocle Bîrsan, Iași

G189. Three spheres are considered with their volumes v_1, v_2, v_3 , their areas s_1, s_2, s_3 and the lengths of their equatorial circles l_1, l_2, l_3 . Prove that

$$\frac{v_1}{l_1 + l_2} + \frac{v_2}{l_2 + l_3} + \frac{v_3}{l_3 + l_1} \geq \frac{1}{12\pi}(s_1 + s_2 + s_3).$$

D.M. Bătinețu-Giurgiu, București

G190. If x and y are irrational numbers such that $x^2 + y$, $y^2 + x$ and $x + y$ are rational, prove that $xy < \frac{1}{4}$.

Neculai Moraru, Suceava and Silviu Boga, Iași

G191. Let $A = \{1, 2, \dots, n\}$ with $n \in \mathbb{N}, n \geq 5$. Prove that the set A can be written as the union of two disjoint sets B and C so that the product of the elements of B be equal to the sum of elements of C .

Gheorghe Iurea, Iași

G192. Prove that, in any non-isosceles triangle ABC , six points $M, N \in [BC]$, $P, Q \in [AB]$ and $R, S \in [AC]$ exist such that the segments $[AU]$, $[BT]$ and $[CV]$ are the sides of an acute triangle, any would be the points $U \in [MN]$, $V \in [PQ]$ and $T \in [RS]$.

Marius Drăgan, București

G193. Effectively using 24 matches, build a polygon with its area equal to 7 units, where a unit area is the area of a square with one match stick side. (*Related to a problem from the book "Mathematical Amusements" by Martin Gardner*).

Dumitru Mihalache, Bârlad

G194. Let $ABCD$ be a convex quadrilateral, and let the half lines $[CK$ and $[CL$ be the angle-bisectors of the angles \widehat{ACD} and respectively \widehat{ACB} ($K \in AD, L \in AB$). Prove that the line LK passes through the centre of the circle inscribed in the triangle ABD if and only if the quadrilateral $ABCD$ can be inscribed into a circle.

Neculai Roman, Mircești (Iași)

G195. The points M and P lie on the non-parallel side (AD) of the trapezoid $ABCD$, such that $M \in [AP]$. The parallel lines to the two bases intersect BC at N and at Q , respectively. Prove that the trapezia $ABNM$ and $PQCD$ have their diagonals respectively parallel if and only if CD and PQ are directly proportional to MN and AB (*Connected with problem G105 of RecMat 1/2006*).

Claudiu Ștefan Popa, Iași

B. Highschool Level

L186. Determine the measures of the angles of triangle ABC knowing that

$$\operatorname{tg} A + 2\sqrt{\operatorname{tg} B} + 3\sqrt[3]{\operatorname{tg} C} = 6\sqrt[6]{\operatorname{tg} A + \operatorname{tg} B + \operatorname{tg} C}.$$

Cătălin Calistru, Iași

L187. Let ABC be a triangle, and let the points $D, L \in (BC)$, $E, M \in (CA)$ and $F, N \in (AB)$. Denote $\{P\} = FE \cap AL$, $\{Q\} = DF \cap BM$, $\{S\} = ED \cap CN$. Prove that if two of the assertions below hold true the third one is also true:

- (i) the straight lines AD, BE and CF are concurrent;
- (ii) the straight lines AL, BM and CN are concurrent;
- (iii) the straight lines DP, EQ and FS are concurrent.

Titu Zvonaru, Comănești

L188. Let Oxy be a Cartesian system of coordinates and consider the line $d : x = 1$ with the point $M \in d$. The point P is obtained from M by a rotation of centre O and angle 2α , where α is the directed angle \widehat{yOM} . Write the equation $y = f(x)$ of

the curve described by P when M runs along the line d and draw the graph of this curve.

Temistocle Bîrsan, Iași

L189. A system of $n \geq 3$ coplanar points A_1, A_2, \dots, A_n , with any three among them being not collinear, is said to be *good* when for any $1 \leq i < j < k \leq n$ and the points $M \in [A_i A_j]$, $N \in [A_j A_k]$ and $P \in [A_k A_i]$, the inequality $MN^2 + NP^2 + MP^2 \leq A_i A_j^2 + A_j A_k^2 + A_k A_i^2$ holds. Determine the good systems with a maximum number of points.

Vlad Emanuel, student, București

L190. Prove that

$$8\sqrt{15} \leq (\sqrt{x} + \sqrt{x+48})(\sqrt{16-x} + \sqrt{1-x}) \leq 36, \quad \forall x \in [0, 1].$$

Gheorghe Iurea, Iași

L191. Show that any three positive real numbers x, y, z satisfy the inequality

$$\begin{aligned} (x^2 + y^2 + z^2) \left(\frac{1}{(x+y)(x+z)} + \frac{1}{(x+y)(y+z)} + \frac{1}{(x+z)(y+z)} \right) &\geq \frac{9}{4} \geq \\ &\geq (xy + xz + yz) \left(\frac{1}{(x+y)(x+z)} + \frac{1}{(x+y)(y+z)} + \frac{1}{(x+z)(y+z)} \right). \end{aligned}$$

Marian Tetiva, Bârlad

L192. Let \mathcal{P} be the set of monic polynomials with their coefficients in the interval $[0, 9]$. Determine the supremum of the modules of the complex roots of polynomials in \mathcal{P} .

Adrian Reisner, Paris

L193. Let $A \in M_3(\mathbb{R})$ be a symmetric matrix with its trace equal to its determinant. Prove that $2 \cdot \text{Tr}(A^{n+2}) \geq \text{Tr} A \cdot (\text{Tr}(A^{n+1}) - \text{Tr}(A^{n-1}))$, $\forall n \in \mathbb{N}^*$.

Paul Georgescu and Gabriel Popa, Iași

L194. Let $p \geq 2$ be a natural number and consider the equation $\frac{1}{\sqrt[p]{1+x}} + \frac{1}{\sqrt[p]{2+x}} + \dots + \frac{1}{\sqrt[p]{n+x}} = \sqrt[p]{x^{p-1}}$. Prove that, for any $n \in \mathbb{N}^*$, this equation has a unique positive solution x_n , show that the sequence $(x_n)_{n \geq 1}$ is monotonic and calculate the limit $\lim_{n \rightarrow \infty} \frac{n}{x_n}$.

Gabriel Dospinescu, student, Paris

L195. Let $(a_n)_{n \geq 1}$ be a sequence of real numbers that are greater than 1, such that $a_{n+1} \geq 2a_n - 1$, $\forall n \in \mathbb{N}^*$. We define $x_n = \frac{1}{a_1} + \frac{1}{a_1 a_2} + \dots + \frac{1}{a_1 a_2 \dots a_n}$, $\forall n \geq 1$.

a) Prove that the sequence $(x_n)_{n \geq 1}$ converges (and denote by x its limit).

b) Show that the sequence $(a_n)_{n \geq 1}$, where $x = x_n + \frac{\alpha_n}{a_1 a_2 \dots a_n (a_{n+1} - 1)}$, $\forall n \geq 1$, is monotonic and convergent, and determine its limit.

Dumitru Mihalache and Marian Tetiva, Bârlad